SMITHSONIAN SCIENCE INFOR Project number (Do <b>not</b> us	MATION EXCHANGE e this space)	U.S. DEPARTMI HEALTH, EDUCATION PUBLIC HEALTI NOTICE	, AND WELFARE I SERVICE OF	PROJECT NUMBER  ZO1 HL 00011-02 LBG	
	İ	INTRAMURAL RESEA	CH PROJECT		
PERIOD COVERED July 1, 1976 thr	ough Septemb	er 30, 1977			
TITLE OF PROJECT (80 char	acters or less)				
The Development	of Chick Emb	ryo Retina			
NAMES, LABORATORY AND INS PROFESSIONAL PERSONNEL EN	NGAGED ON THE PR		OF PRINCIPAL IN	IVESTIGATORS AND ALL OTHER	
PI: F. G. De Marshall	Mello Nirenberg	Visiting A Chief, LBG		LBG NHLBI LBG NHLBI	
COOPERATING UNITS (if any	<sub>Y</sub> )				
LAB/BRANCH					
Laboratory of Bi	ochemical Ge	netics			
Section on Molec	ular Riology	•		,	
INSTITUTE AND LOCATION	dial biology				
NHLBI, NIH, Beth	ondo Marvila	ind 20014			
TOTAL MANYEARS:	PROFESSION	<del></del>	OTHER:		
0.5	0.5		ornen:		1
CHECK APPROPRIATE BOX(ES)		<del></del>	1		
□ (a) HUMAN SUBJECTS	•	) HUMAN TISSUES	Γ¥	(c) NEITHER	
_ (,	Ц (	, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	( <u>4</u> )	(c) nermen	
☐ (a1) MINORS ☐ (a2)	NTERVIEWS				
SUMMARY OF WORK (200 work	ds or less - und	erline keywords)			
Come Lineba				. differentiation	
				a differentiation were	
				vas characterized in the cine to GABA. 2) Glutam	
					<u></u>
acid decarboxylase specific activity in the retina, increased during the course of embryonic development, either when measured in ovo or in aggregate					
					_
cultures. 3) The presence of GABA in the culture medium prevented the development of glutamic acid decarboxylase activity in aggregate cultures.					
4) The proposed course of this project is to attempt to correlate the					
biochemical chan					
· · · · · · · · · · · · · · · · · · ·	<del>-</del>				Ì
					ļ

## Project Description:

Objectives: The objective of the project is to study the biochemical step required for synaptogenesis in chick embryo retina.

Major Findings: Two pathways for  $\gamma$ -aminobutyric acid synthesis were found in chick embryo retina. The first pathway depends upon the conversion of putrescine to ornithine decarboxylase and the subsequent conversion of ornithine to  $\gamma$ -aminobutyric acid. The second route of synthesis is dependent upon the conversion of glutamic acid to  $\gamma$ -aminobutyric acid, catalyzed by glutamic acid decarboxylase. Elevation of cAMP levels in neuroblastoma cells was shown to induce ornithine decarboxylase activity. Thus, in the developing embryo, neurotransmitters which affect cAMP levels may regulate ornithine decarboxylase activity and thereby control the rate of GABA synthesis from ornithine.

GABA was found to regulate the specific activity of glutamic acid decarboxylase in cells dissociated from chick embryo retina and cultured in vitro.

Significance to Biomedical Research: These findings show that GABA can be synthesized by a novel pathway and provide new insight on the relationship and regulation of GABA synthesis.

Proposed Course: To determine the effect of retina neurotransmitters and other compounds on retina synaptogenesis.

## Publications:

1. De Mello, F. G., Bachrach, U. and Nirenberg, M.: Ornithine and glutamic acid decarboxylase activities in the developing retina. J. Neurochem. 27: 847-851, 1976.